

PHYS 3344

Fall 2019

TE Coan

Due: 27 Sep '17 6:00 pm

Homework 3

1. Even though the total force on a system of N particles ($\sum_{\alpha} \mathbf{F}_{\alpha}^{\text{ext}} + \sum_{\alpha} \sum_{\beta \neq \alpha} \mathbf{F}_{\alpha\beta}$) is zero, the net torque may not be zero. Show that the net torque has the same value in any coordinate system. **Hint:** Calculate the net torque on a system of particles in any two fixed coordinate systems whose origins differ by a fixed vector amount \mathbf{a} .

2. The speed v of some strangely shaped projectile of mass m varies with distance x as $v(x) = \alpha x^{-n}$. Assume that at $t = 0$, $v(x = 0) = 0$.

a) What force $F(x)$ is responsible for this state of affairs?

b) What is $x(t)$?

c) Finally, what is $F(t)$?

3. A DART train moves along the tracks at a constant speed u . A woman on the train throws a ball, for reasons mysterious to this day, of mass m straight ahead with a speed v with respect to *herself*.

a) What is the kinetic energy gain ΔKE_{train} of the ball as measured by a person on the train?


b) Compute the same quantity as in (a) but for a person standing by the *track*?

c) How much work W_{Woman} is done by the woman throwing the ball?

d) How much Work W_{Train} is done by the train?

4. A chain of length b and mass ρb is suspended from one end at a height b above a table so that the free end barely touches the tabletop. At time $t = 0$, the fixed end of the chain is released. Find the force that the tabletop exerts on the chain after the original fixed end has fallen a vertical distance x .

5. This is a somewhat involved problem, but fully within your ability to solve. It is designed to show you the advantages of a multi-stage rocket compared to a single-stage rocket. Since the commercialization of space by Mr. Amazon and Mr. Tesla seems approaching, it seems sensible to know something about it. Suppose that the payload (e.g., a capsule carrying some hyper-affluent people, or maybe something more interesting) has a mass m and is mounted on a two-stage rocket (see the fig.). The *total* mass –both rocket stages fully fueled, plus the payload – is Nm . The mass of the second-stage plus the payload, after first-stage burnout and separation, is nm . In each case, the ratio of the burnout mass (casing) to initial mass (casing plus fuel) is r , and the exhaust speed of the gas with respect to either stage is v_{ex} .



a) Show that the velocity v_1 gained after the first stage burn, starting from rest (and ignoring gravity), is given by

$$v_1 = v_{ex} \ln \left[\frac{N}{rN + n(1 - r)} \right]$$

b) Obtain a corresponding expression for the additional velocity v_2 gained from the second-stage burn. Box your answer.

c) Adding v_1 and v_2 , you have the payload velocity v in terms of N , n , and r . Taking N and r as constants, find the value of n that maximizes v . **HINT:** This will turn out to be a simple function of N . The algebra can seem messy so I recommend NOT combining the natural logs when you go to maximize v . You can also use Mathematica. Box your answer.

d) Show that the condition for v to be a maximum corresponds to having equal velocity gains for each of the two stages. Find the maximum value of v . Comment on why this expression makes sense. Box your answer.

6. A particle of mass m at the end of a light string wraps itself about a fixed vertical cylinder of radius a (see the figure below). All the motion is in the horizontal plane (ignore gravity). The angular velocity of the string is ω_0 when the distance from the particle to the point of contact of the string and cylinder is b . Find the angular velocity and tension T in the string after the the string has turned through an additional angle θ .

