


July 8, 2015, Pittsburgh

Jets

Zoltan Nagy
DESY


What are Jets?

A di-jet ATLAS event




What are Jets?

A multi-jet (6-jet) event




What are Jets?

- The p_T is concentrated in a few narrow sprays of particles
- These sprays are called jets.
- Events with big total p_T are rather rare...
- ... but when they happen, the p_T is always in jets



Why are the Jets there?

Here is a Feynman graph for quad-quark scattering with additional radiation that can contribute to the jet events.



- Initial state


- If $p_1 \rightarrow 0$, then $1/(p_a - p_1)^2 \rightarrow \infty$
- If $p_1 \rightarrow \lambda p_a$, then $1/(p_a - p_1)^2 \rightarrow \infty$

- Final state

- If $p_2 \rightarrow 0$, then $1/(p_2 + p_3)^2 \rightarrow \infty$
- If $p_3 \rightarrow 0$, then $1/(p_2 + p_3)^2 \rightarrow \infty$
- If $p_3 \rightarrow \lambda p_2$, then $1/(p_2 + p_3)^2 \rightarrow \infty$

The probability is big to get a spray of collimated particles plus some low momentum particles with wide angle.

How many Jets are there?






ATLAS
EXPERIMENT


Run Number: 158548, Event Number: 2486978

Date: 2010-07-04 06:46:45 CEST

**Multijet Event in
7 TeV Collisions**



How many Jets are there?






ATLAS EXPERIMENT

Run Number: 158548, Event Number: 2486978

Date: 2010-07-04 06:46:45 CEST

Multijet Event in 7 TeV Collisions




How many Jets are there?

Jet structure at large resolution scale:

The jet algorithm find one fat jet

These hadrons are part of the "beam jet" when the jet resolution is crude.




How many Jets are there?

Jet structure at small resolution scale:

The jet algorithm find one fat jet

Now, they are resolved as a jet.

These are still part of the beam jet.




The number of the jets depend on the *typical resolution scale (theory)*, *detector sensitivity and angular resolution (experiment)* .

Jets in pQCD

- Let us consider a 3-jet event in e^+e^- annihilation with the typical resolution scale Q .
- At each vertex in a diagram, there is a **factor of the strong coupling**, $g_s^2/(4\pi) = \alpha_s$
- The simplest graph that contributes to this process is the tree level graph

Tree level graph




*All the three **patrons** are well separated from each others and the “distance” is measured by some hardness variable like transverse momentum or virtuality.*



Jets in pQCD

- In the perturbation theory should consider radiative correction.
- We can consider **one more gluon** in the final state...

Resolvable real radiation



Unresolvable real radiation




4-jet configuration
all the four partons are well separated


Jets in pQCD

- In the perturbation theory should consider radiative correction.
- We can consider **one more gluon** in the final state...

Resolvable real radiation



Unresolvable real radiation




4-jet configuration

all the four partons are well separated


Jets in pQCD

- In the perturbation theory should consider radiative correction.
- We can consider **one more gluon** in the final state...

Resolvable real radiation



Unresolvable real radiation




- Everything inside the green cone is unresolvable and integrated out.
- It is a **singular integral**.
- This singularity **has to be cancelled**. Otherwise we cannot make pQCD predictions for jet production.

4-jet configuration


all the four partons are well separated

Jet in pQCD


We have to also consider the **virtual corrections**, thus we have graphs like...



Approximately we can omit the real and loop graph by **adjusting** $\alpha_s(Q^2)$ and the "normalization" of the tree graph $Z(Q^2)$.



- Singularities has to be cancelled between the two graphs!!!
- This cancelation has to be ensured by the jet definition!!!




INFRARED SAFETY

Infrared Safety

The jet algorithm has to be infrared safe. This means it has to be *insensitive for any small scale physics* (soft or collinear radiation).

- We construct jets from particle momenta $\{p_1, p_2, \dots, p_m\}$.
- We get N jets with momenta $\{P_1, P_2, \dots, P_N\}$.



- If any p_i becomes **very small**, we should get the same jets by leaving particle i out.
- If any two momenta p_i and p_j become **collinear**, we should get the same jets by replacing the particles by one with momentum $p_i + p_j$.

Jet Cross Sections

In the general case the cross section is given by

$$\sigma[F] = \sum_m \frac{1}{m!} \int d\{p, f\}_m |M(\{p, f\}_m)|^2 \underbrace{F(\{p\}_m)}_{\text{Jet measurement function}}$$

$F(\{p\}_m) \equiv F(p_1, p_2, \dots, p_m)$

INFRARED SAFETY (formal definition):

$$F(p_1, p_2, \dots, p_m, p_{m+1}) \xrightarrow{p_{m+1} \rightarrow 0} F(p_1, p_2, \dots, p_m)$$


$$F(p_1, p_2, \dots, p_m, p_{m+1}) \xrightarrow{p_m \parallel p_{m+1}} F(p_1, p_2, \dots, p_m + p_{m+1})$$

One can consider for example the inclusive one jet cross section

$$\sigma[F] \implies \frac{d\sigma}{dp_T dy} \quad F(\{p\}_m) \implies \delta(p_T - \underbrace{P_T(\{p\}_m)}_{\text{Transverse momentum of the observed jet}}) \delta(y - \underbrace{Y(\{p\}_m)}_{\text{Rapidity of the observed jet}})$$


One Jet Inclusive Cross Section

A result from ATLAS



- Note nine order of magnitude variation in cross section at one $|y|$.
- Compared to NLO pQCD prediction and the agreement is very good! (No sign of new physics...)

Now, it is high time to give a definition of the jet algorithm.



Jet Algorithms

- There are two kind of algorithms for defining jets:
 - cone algorithms
 - successive combination algorithms
- Both can be infrared safe.
- I will discuss just the successive combination algorithms.
- This traces back to the JADE collaboration at DESY.

THE KT JET ALGORITHM

- Choose an angular resolution parameter R
- Start with the list of protojets, specified by their momenta $\{p_1, p_2, \dots, p_m\}$.
- Start with an empty list of finished jets, $\{\}$.
- The result is a list of finished jets with their momenta, $\{P_1, P_2, \dots, P_N\}$.
- Many are low p_T debris, just ignore them.

kT Jet Algorithm

1. For each pair of protojets define

$$d_{ij} = \min \{p_{T,i}^2, p_{T,j}^2\} [(\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2] / R^2$$

and for each protojet define

$$d_i = p_{T,i}^2$$

2. Find the smallest of the d_{ij} and the d_i

$$d_{\min} = \min_{i,j} \{d_i, d_{ij}\}$$

3. If d_{\min} is a d_{ij} , merge protojets i and j into a new protojets k with momentum


$$p_k = p_i + p_j$$

4. If d_{\min} is a d_i , then protojet i is "not mergable". Remove it from the list of protojets and add it to the list of finished jets.
5. If protojets remain, go to step 1.

kT Jet Algorithm

Why the name?

$$d_{ij} = \min \{p_{T,i}^2, p_{T,j}^2\} [(\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2] / R^2 \approx k_{\perp}^2 / R^2$$



Infrared safety of this:

- Suppose $p_j \rightarrow 0$
 - Then when it merges with other protojet,

$$p_k = p_i + p_j \rightarrow p_i$$


- If it never merges with other protojets, then it just remains as a low p_T jets at the end.

- Suppose $p_i = \lambda p_j$
 - Then protojets i and j are always merged at the beginning to

$$p_k = p_i + p_j$$

Example with kT Algorithm

Here is an event from Cacciari, Salam and Soyes (2008). An event was generated by HERWIG++ along with (lots of) random soft particles.




- The detector area that goes into each jet is **irregular**.
- The k_T algorithm has the tendency to “suck” in low p_T radiation and contaminate the jets with underlying event.

Cambridge-Aachen Algorithm

This is a variation on the general successive combination algorithm. The only difference is in the “distance” measure.

$$d_{ij} = \left[(y_i - y_j)^2 + (\phi_i - \phi_j)^2 \right] / R^2$$
$$d_i = 1$$

Only the angles count!




With this algorithm the jets still have irregular shape.

Anti-kT Algorithm

This is another variation on the general successive combination algorithm. The only difference is in the “distance” measure.

$$d_{ij} = \min \left\{ \frac{1}{p_{T,i}^2}, \frac{1}{p_{T,j}^2} \right\} [(y_i - y_j)^2 + (\phi_i - \phi_j)^2] / R^2$$
$$d_i = \frac{1}{p_{T,i}^2}$$




The highest p_T protojet has the priority to absorb nearby softer protojets.

The high p_T jets are round.

When Fixed Order Brakes Down

Let us consider **2photon + 1jet inclusive** production and plot the di-photon pT distribution

$p p \rightarrow \gamma\gamma + \text{jet}, E_{\text{cm}} = 14 \text{ TeV}$



- For this distribution the characteristic scale is

$$Q^2 = (p_{\gamma\gamma,\perp} - 40 \text{ GeV})^2$$


- The NLO distribution has **discontinuity at 40 GeV**. It is $-\infty$ from the right and $+\infty$ from the left.
- The singularities are logarithms (it appears finite because of the bin smearing effect).
- The effective expansion variable is

$$\alpha_s(Q^2) \log^2 \frac{Q^2}{(40 \text{ GeV})^2}$$

- This effect has to be summed up all order. NLO calculation is not enough.

Jet in pQCD



We have to also consider the **virtual corrections**, thus we have graphs like...




When Q gets small the coupling and the logarithm blow up.

Jet in pQCD

We have to also consider



We can omit top graph by $q^2 > Q^2$ and the bottom graph by $q^2 < Q^2$



logarithm blow up.

Conclusions

- QCD gives us **jets**.
- Jets are **real and seen** in experiments.
- To measure jet cross sections, you need a careful definition of jets.
- At LHC we use successive combination algorithms, such as kT, Cambridge-Aache or **anti-kT algorithm**.
- The definition **needs to be infrared safe**.
- Infrared safety allow us to make pQCD prediction.
 - Fixed order calculations, LO, NLO or NNLO
- Jet cross sections (in general pQCD cross sections) **usually suffers on large logarithms** and these logarithms need to be summed up all order.
 - Summing up logarithms **analytically**
 - Summing up logarithm **numerically by parton shower** algorithms.