

# Lecture 2: Quantum Mechanics and Relativity

## *Atom*

<i>Atomic number</i>	<i>A</i>
<i>Number of protons</i>	<i>Z</i>
<i>Number of neutrons</i>	<i>A-Z</i>
<i>Number of electrons</i>	<i>Z</i>


*Charge of electron = charge of proton  $\sim 1.6 \times 10^{-19} \text{ C}$*


<i>Size of the atom</i>	<i><math>\sim 10^{-10} \text{ m}</math></i>
<i>Size of the nucleus</i>	<i><math>\sim 10^{-15} \text{ m}</math></i>

## Two questions:


- Why did Rutherford need  $\alpha$  – particles to discover the atomic nucleus?
- Why do we need huge accelerators to study particle physics today?

## Observation of very small objects using visible light








Aperture diameter:  $D = 20\mu\text{m}$   
Focal length: 20 cm



**Observation of light diffraction, interpreted as evidence that light consists of waves since the end of the 17<sup>th</sup> century**

**Angular aperture of the first circle (before focusing):**

$$\alpha = 1.22\lambda / D$$




Opaque disk, diam.  $10\mu\text{m}$   
in the center


**Presence of opaque disk is detectable**

# Opaque disk of variable diameter


diameter =  $.4 \mu\text{m}$




diameter =  $.2 \mu\text{m}$



diameter =  $.1 \mu\text{m}$



no opaque disk




The presence of the opaque disk in the center is detectable if its diameter is larger than the wavelength  $\lambda$  of the light

The RESOLVING POWER of the observation depends on the wavelength  $\lambda$

Visible light: not enough resolution to see objects smaller than  $0.2 - 0.3 \mu\text{m}$

# Opaque screen with two circular apertures



aperture diameter:  $10\ \mu\text{m}$   
distance between centers:  $15\ \mu\text{m}$





Image obtained by shutting one aperture alternatively for 50% of the exposure time





Image obtained with both apertures open simultaneously

Light is a wave and can interfere !

# Photoelectric effect: evidence that light consists of particles



**Observation of a threshold effect as a function of the frequency (wavelength) of the light impinging onto the electrode at negative voltage (cathode):**

**Frequency  $\nu < \nu_0$  : electric current = zero, independent of luminous flux;**


**Frequency  $\nu > \nu_0$  : current  $> 0$ , proportional to luminous flux**

**INTERPRETATION (A. Einstein):**

- Light consists of particles (“photons”) !!!!
- Photon energy proportional to frequency:

$$E = h \nu \quad (\text{Planck constant } h = 6.626 \times 10^{-34} \text{ J s})$$

- Threshold energy  $E_0 = h\nu_0$ : the energy needed to extract an electron from an atom (depends on the cathode material)




Albert Einstein

Repeat the experiment with two circular apertures using a very weak light source

**Luminous flux = 1 photon /second**

(detectable using modern, commercially available photomultiplier tubes)

**Need very long exposure time**




aperture diameter:  $10\ \mu\text{m}$   
distance between centres:  $15\ \mu\text{m}$

Question: which aperture will photons choose?

Answer: diffraction pattern corresponds to both apertures simultaneously open, independent of luminous flux

*Interference pattern*



**Photons have both particle and wave properties simultaneously**

**It is impossible to know which aperture the photon traversed**

**The photon can be described as a coherent superposition of two states**

# Black body radiation

Electromagnetic radiation emitted by hot object.

Statistical mechanics lead to “ultraviolet catastrophe”

i.e., amount of energy emitted at short wavelength became infinite.

1900 Planck: One can avoid the UV catastrophe and describe the experimentally measured spectrum IF electromagnetic radiation is Quantized, i.e. comes in little packages of energy


$$E = h\nu$$

Planck's constant

$$h = 6.626 \times 10^{-27} \text{ ergs}$$

$\nu$  - frequency

$\lambda$  - wavelength  $\sim 1/\nu$





## 1924: De Broglie's principle

**Not only light, but also matter particles possess both the properties of waves and particles**

**Relation between wavelength and momentum:**

$$\lambda = \frac{h}{p}$$

$h$ : Planck constant

$p = m v$  : particle momentum



Louis de Broglie

**Hypothesis soon confirmed by the observation of diffraction pattern in the scattering of electrons from crystals, confirming the wave behaviour of electrons** (Davisson and Germer, 1927)

**Wavelength of the  $\alpha$  – particles used by Rutherford in the discovery of the atomic nucleus:**

$$\lambda = \frac{h}{m_{\alpha} v} \approx \frac{6.626 \times 10^{-34} \text{ J s}}{(6.6 \times 10^{-27} \text{ kg}) \times (1.5 \times 10^7 \text{ m s}^{-1})} \approx 6.7 \times 10^{-15} \text{ m} = 6.7 \times 10^{-13} \text{ cm}$$

$\alpha$  particle mass      0.05 c      ~ resolving power of Rutherford's experiment

## Typical tools to study objects of very small dimensions

		<b>Resolving power</b>
<b>Optical microscopes</b>	<b>Visible light</b>	$\sim 10^{-6}$ m
<b>Electron microscopes</b>	<b>Low energy electrons</b>	$\sim 10^{-9}$ m
<b>Radioactive sources</b>	<b><math>\alpha</math>-particles</b>	$\sim 10^{-14}$ m
<b>Accelerators</b>	<b>High energy electrons, protons</b>	$\sim 10^{-18}$ m

## Units in Particle Physics

**Fundamental Units in Physics:** **mass, length, time** (**m, kg, s**) are not very useful in the world of particle physics.

**Typical dimensions are:**

**Size of the nucleus**  $\sim 1 \text{ fermi} = 10^{-15} \text{ m}$

**Mass of the proton**  $m_p = 1.672 \cdot 10^{-27} \text{ kg}$

### Energy


#### **1 electron-Volt (eV):**

the energy of a particle with electric charge =  $|e|$ , initially at rest, after acceleration by a difference of electrostatic potential = 1 Volt

( $e = 1.60 \times 10^{-19} \text{ C}$ )

$$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$$

$$\text{J} = \text{kg m/s}^2$$



## Multiples:

deca	da	$10^1$	deci	d	$10^{-1}$
hecto	h	$10^2$	centi	c	$10^{-2}$
kilo	k	$10^3$	mili	m	$10^{-3}$
mega	M	$10^6$	micro	$\mu$	$10^{-6}$
giga	G	$10^9$	nano	n	$10^{-9}$
tera	T	$10^{12}$	pico	p	$10^{-12}$
peta	P	$10^{15}$	femto	f	$10^{-15}$

### Energy:

$$1 \text{ keV} = 10^3 \text{ eV} ; \quad 1 \text{ MeV} = 10^6 \text{ eV}$$

$$1 \text{ GeV} = 10^9 \text{ eV}; \quad 1 \text{ TeV} = 10^{12} \text{ eV}$$

**Energy of a proton in the LHC (in the year 2007):  $7 \text{ TeV} = 1.12 \times 10^{-6} \text{ J}$**

**This energy is equal to a body of mass = 1 mg moving at speed = 1.5 m /s (a bee)**

**The conversion constant between MKS and particle physics units is**

$$\mathbf{hc = 197.327 \text{ MeV fm}}$$

**My rest mass (weight =80 kg) is:**

$$\mathbf{M = 80 \text{ kg} \cdot (3 \cdot 10^8 \text{ m/s})^2 / 1.6 \cdot 10^{-19} \text{ J/eV} = 45 \cdot 10^{14} \text{ TeV/c}^2}$$

# Energy and momentum for relativistic particles

Speed of light in vacuum  $c = 2.99792 \times 10^8$  m / s (in the absence of magnetic field)

**Total energy:** 
$$E = mc^2 = \frac{m_0 c^2}{\sqrt{1 - (v/c)^2}}$$
  $\left( \begin{array}{l} m: \text{ relativistic mass} \\ m_0: \text{ rest mass} \end{array} \right)$

**Expansion in powers of  $(v/c)$ :** 
$$E = m_0 c^2 + \frac{1}{2} m_0 v^2 + \dots$$
  
(only valid for  $v/c$  small)

↑  
energy  
associated  
with rest mass


↑  
“classical”  
kinetic  
energy

**Momentum:** 
$$p = mv = \frac{m_0 v}{\sqrt{1 - (v/c)^2}}$$

$$\frac{pc}{E} = \frac{v}{c} \equiv \beta$$

$E^2 - p^2c^2 = (m_0c^2)^2$  “relativistic invariant” (effective mass)  
(same value in all reference frames)

Special case: the photon ( $v = c$  in vacuum)

Einstein  $E = h \nu$   
de Broglie  $\lambda = h / p$    $E / p = v \lambda = c$  (in vacuum)  
 $E^2 - p^2c^2 = 0$   
photon rest mass  $m_g = 0$

Momentum units: eV/c (or MeV/c, GeV/c, ...)

Mass units: eV/c<sup>2</sup> (or MeV/c<sup>2</sup>, GeV/c<sup>2</sup>, ...)

Numerical example: electron with  $v = 0.99 c$

Rest mass:  $m_e = 0.511 \text{ MeV}/c^2$

$\gamma \equiv \frac{1}{\sqrt{1 - (v/c)^2}} = 7.089$  (often called “Lorentz factor”)

Total energy:  $E = \gamma m_e c^2 = 7.089 \times 0.511 = 3.62 \text{ MeV}$

Momentum:  $p = (v / c) \times (E / c) = 0.99 \times 3.62 = 3.58 \text{ MeV}/c$

## **“Natural System of Units”**

$$c = 1$$

$$E^2 = p^2 + m^2$$

- **All quantities in multiples of eV**

- **Unit of length**                      **1 fm = 1/197.3 MeV**

- **Unit of time**                        **1 s = 1/6.58 10<sup>-22</sup> MeV**