
4321

1. Simplify these expressions:

- (a) $\int_{x=-\infty}^{+\infty} \delta(x) \sin(x) dx$
- (b) $\int_{x=-\infty}^{+\infty} \delta(x) \sin(t) dx$
- (c) $\int_{x=-\infty}^{+\infty} \delta(x - t) \sin(x) dx$
- (d) $\int_{x=-\infty}^{+\infty} \delta(x - t) \sin(x + t) dx$
- (e) $\int_{x=-\infty}^{+\infty} \delta(3x) \cos(x) dx$
- (f) $\int_{x=14}^{+\infty} \delta(3x) \cos(x) dx$
- (g) $\int_{x=-\infty}^{+\infty} \delta'(x)(3x^2 + x + 7) dx$, where the prime means derivative.
- (h) $\int_{x=-\infty}^{+\infty} \delta''(x)(3x^2 + x + 7) dx$
- (i) $\int_{x=-\infty}^{+\infty} \delta''(x + 2)(3x^2 + x + 7) dx$

2. Use the Heavyside or step function $\theta(t)$ to define the graph of a square wave pulse of height 4 that begins at $t=3$ and ends at $t=6$ using:

- (a) two θ functions additively.
- (b) two θ functions multiplicatively.
- (c) a single θ function.

3. Sketch the graph of $\int_{x'=-\infty}^x \left[\int_{x''=-\infty}^{x'} \delta(x'') dx'' \right] dx'$ vs. x .

7305

1. If a and b are constants and $\delta^{(m)}$ means the m 'th derivative of the delta function, show that $x^n \delta^{(m)}(x) =$

- (a) 0, if $m < n$
- (b) $(-1)^n n! \delta(x)$, if $m = n$
- (c) $\frac{(-1)^n m!}{(m-n)!} \delta^{(m-n)}(x)$, if $m > n$

Don't forget that these distributions only make physical sense inside an integral and multiplied by a test function $f(x)$ with bounded support.

2. You are most familiar with the Euclidean norm of an n -dimensional vector $\vec{r} = \sum_{i=1}^n x_i \hat{e}_i$,

$$|\vec{r}| = \sqrt{\sum_{i=1}^n (x_i)^2} = \left(\sum_{i=1}^n |x_i|^2 \right)^{\frac{1}{2}}$$

that assigns to every non-zero vector a strictly positive length, but this only one possible norm. In general the p -norm is

$$||\vec{r}||_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}$$

- (a) Explain why $p = 1$ is called the “taxicab norm”.
- (b) Explain why $p \rightarrow \infty$ is called the “maximum norm”.
- (c) Draw graphs of “circles” in $n=2$ dimensions (that is, in the x,y -plane) of unit norm $\|\vec{r}\|_p = 1$ for
 - i. $p = 1$
 - ii. $p = 1.5$
 - iii. $p = 2$ (This is the only true circle.)
 - iv. $p = 3$
 - v. $p = 6$
 - vi. $p \rightarrow \infty$

and describe what happens to the shape as p increases. What happens for $p < 1$?

Bonus: Solve as much of the other class’ assignment as you can.