1. In lecture, we claimed that the error

$$E_N(c_1,\ldots,c_N) \equiv \int_a^b d\xi \ |f(\xi) - f_N(\xi)|^2 = \int_a^b d\xi \ [f(\xi) - f_N(\xi)]^* [f(\xi) - f_N(\xi)]$$

made in the approximation

$$f_N(\xi) = \sum_{n=1}^N c_n u_n(\xi)$$

to the function $f(\xi)$ was minimized if the expansion coefficients were chosen as

$$c_n = \int_a^b d\xi \ u_n^*(\xi) f(\xi) \quad .$$

Prove this assertion. Hint: Write $c_n = a_n + ib_n$ where a_n and b_n are real constants, and $\int_a^b d\xi \ u_n^*(\xi) f(\xi) = A_n + iB_n$ where A_n and B_n are real constants. Then require that

$$\frac{\partial E_N}{\partial a_k} = 0 = \frac{\partial E_N}{\partial b_k}$$

2. Find the potential $\Phi(x, y, z)$ inside a parallelopiped (0 < x < a, 0 < y < b, 0 < z < c) for which the x-component of the electric field is specified on the face at x = 0, and the others faces are grounded.

$$E_x(0, y, z) = f(y, z)$$

$$\Phi(a, y, z) = 0 = \Phi(x, 0, z) = \Phi(x, b, z) = \Phi(x, y, 0) = \Phi(x, y, c)$$

Bonus:



Two perfectly conducting infinite and grounded half-planes intersect at an angle $\pi/3$ radians. A charge q is located halfway between the planes at a distance d from the intersection. Find the potential energy of the configuration.