

PHYS 1303 Final Exam Example Questions

1. Which quantity can be converted from the English system to the metric system by the conversion factor

$$\frac{5280 \text{ f}}{\text{mi}} \cdot \frac{12 \text{ in}}{\text{f}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{1 \text{ h}}{3600 \text{ s}} ?$$

- a. feet per second
 - b. feet per hour
 - c. miles per second
 - d. miles per hour
 - e. miles per minute
2. A particle moving with a constant acceleration has a velocity of 20 cm/s when its position is $x = 10 \text{ cm}$. Its position 7.0 s later is $x = -30 \text{ cm}$. What is the acceleration of the particle?
- a. -7.3 cm/s^2
 - b. -8.9 cm/s^2
 - c. -11 cm/s^2
 - d. -15 cm/s^2
 - e. -13 cm/s^2
3. A rocket, initially at rest, is fired vertically with an upward acceleration of 10 m/s^2 . At an altitude of 0.50 km, the engine of the rocket cuts off. What is the maximum altitude it achieves?
- a. 1.9 km
 - b. 1.3 km
 - c. 1.6 km
 - d. 1.0 km
 - e. 2.1 km
4. In a location where the train tracks run parallel to a road, a high speed train traveling at 60 m/s passes a car traveling at 30 m/s in the opposite direction. How long does it take for the train to be 180 m away from the car?
- a. 2.0 s
 - b. 3.0 s
 - c. 6.0 s
 - d. 9.0 s
 - e. 18.0 s

5. At $t = 0$, a particle leaves the origin with a velocity of 12 m/s in the positive x direction and moves in the xy plane with a constant acceleration of $(-2.0\hat{i} + 4.0\hat{j})\text{m/s}^2$. At the instant the y coordinate of the particle is 18 m, what is the x coordinate of the particle?

- a. 30 m
- b. 21 m
- c. 27 m
- d. 24 m
- e. 45 m

6. The site from which an airplane takes off is the origin. The x -axis points east; the y -axis points straight up. The position and velocity vectors of the plane at a later time are given by

$$\vec{r} = (1.61 \times 10^4 \hat{i} + 9.00 \times 10^3 \hat{j}) \text{ m} \text{ and } \vec{v} = (150\hat{i} - 21\hat{j}) \frac{\text{m}}{\text{s}}.$$


The magnitude, in meters, of the plane's displacement from the origin is

- a. 9.14×10^3 .
- b. 1.61×10^4 .
- c. 1.84×10^4 .
- d. $9.14 \times 10^3 t$.
- e. $1.61 \times 10^4 t$.

7. A 3.0-kg block slides on a frictionless 20° inclined plane. A force of 16 N acting parallel to the incline and up the incline is applied to the block. What is the acceleration of the block?

- a. 2.0 m/s^2 down the incline
- b. 5.3 m/s^2 up the incline
- c. 2.0 m/s^2 up the incline
- d. 3.9 m/s^2 down the incline
- e. 3.9 m/s^2 up the incline

8. In the figure, if $F = 2.0 \text{ N}$ and $M = 1.0 \text{ kg}$, what is the tension in the connecting string? The pulley and all surfaces are frictionless.



- a. 2.6 N
- b. 1.1 N
- c. 2.1 N
- d. 1.6 N
- e. 3.7 N

9. A 1.0-kg block is pushed up a rough 22° inclined plane by a force of 7.0 N acting parallel to the incline. The acceleration of the block is 1.4 m/s^2 up the incline. Determine the magnitude of the force of friction acting on the block.

- a. 1.9 N
- b. 2.2 N
- c. 1.3 N
- d. 1.6 N
- e. 3.3 N

10. A race car travels 40 m/s around a banked (45° with the horizontal) circular (radius = 0.20 km) track. What is the magnitude of the resultant force on the 80-kg driver of this car?

- a. 0.68 kN
- b. 0.64 kN
- c. 0.72 kN
- d. 0.76 kN
- e. 0.52 kN


11. A 30-kg child rides on a circus Ferris wheel that takes her around a vertical circular path with a radius of 20 m every 22 s. What is the magnitude of the resultant force on the child at the highest point on this trajectory?

- a. 49 N
- b. 0.29 kN
- c. 0.34 kN
- d. 0.25 kN
- e. 0.76 kN

12. A 0.60-kg object is suspended from the ceiling at the end of a 2.0-m string. When pulled to the side and released, it has a speed of 4.0 m/s at the lowest point of its path. What maximum angle does the string make with the vertical as the object swings up?

- a. 61°
- b. 54°
- c. 69°
- d. 77°
- e. 47°

13. A 20-kg mass is fastened to a light spring ($k = 380 \text{ N/m}$) that passes over a pulley as shown. The pulley is frictionless, and the mass is released from rest when the spring is unstretched. After the mass has dropped 0.40 m, what is its speed?



- a. 2.2 m/s
b. 2.5 m/s
c. 1.9 m/s
d. 1.5 m/s
e. 3.6 m/s
14. The only force acting on a 2.0-kg body moving along the x axis is given by $F_x = (2.0x) \text{ N}$, where x is in m. If the velocity of the object at $x = 0$ is $+3.0 \text{ m/s}$, how fast is it moving at $x = 2.0 \text{ m}$?
- a. 4.2 m/s
b. 3.6 m/s
c. 5.0 m/s
d. 5.8 m/s
e. 2.8 m/s
15. An 80-g particle moving with an initial speed of 50 m/s in the positive x direction strikes and sticks to a 60-g particle moving 50 m/s in the positive y direction. How much kinetic energy is lost in this collision?
- a. 96 J
b. 89 J
c. 175 J
d. 86 J
e. 110 J

- 16.A 3.0-kg ball with an initial velocity of $(4\mathbf{i} + 3\mathbf{j})$ m/s collides with a wall and rebounds with a velocity of $(-4\mathbf{i} + 3\mathbf{j})$ m/s. What is the impulse exerted on the ball by the wall?
- $+24\mathbf{i}$ N s
 - $-24\mathbf{i}$ N s
 - $+18\mathbf{j}$ N s
 - $-18\mathbf{j}$ N s
 - $+8.0\mathbf{i}$ N s
17. At the instant a 2.0-kg particle has a velocity of 4.0 m/s in the positive x direction, a 3.0-kg particle has a velocity of 5.0 m/s in the positive y direction. What is the speed of the center of mass of the two-particle system?
- 3.8 m/s
 - 3.4 m/s
 - 5.0 m/s
 - 4.4 m/s
 - 4.6 m/s
- 18.A 4.2-kg object, initially at rest, “explodes” into three objects of equal mass. Two of these are determined to have velocities of equal magnitudes (5.0 m/s) with directions that differ by 90° . How much kinetic energy was released in the explosion?
- 70 J
 - 53 J
 - 60 J
 - 64 J
 - 35 J
- 19.A wheel rotates about a fixed axis with an initial angular velocity of 20 rad/s. During a 5.0-s interval the angular velocity increases to 40 rad/s. Assume that the angular acceleration was constant during the 5.0-s interval. How many revolutions does the wheel turn through during the 5.0-s interval?
- 20 rev
 - 24 rev
 - 32 rev
 - 28 rev
 - 39 rev
- 20.A wheel rotating about a fixed axis with a constant angular acceleration of 2.0 rad/s^2 starts from rest at $t = 0$. The wheel has a diameter of 20 cm. What is the magnitude of the total linear acceleration of a point on the outer edge of the wheel at $t = 0.60 \text{ s}$?
- 0.25 m/s^2
 - 0.50 m/s^2
 - 0.14 m/s^2
 - 0.34 m/s^2
 - 0.20 m/s^2


21. A particle whose mass is 2 kg moves in the xy plane with a constant speed of 3 m/s in the x -direction along the line $y = 5$. What is its angular momentum (in $\text{kg} \cdot \text{m}^2/\text{s}$) relative to the origin?

- a. $-30 \hat{\mathbf{k}}$
- b. $30 \hat{\mathbf{k}}$
- c. $-15 \hat{\mathbf{k}}$
- d. $15 \hat{\mathbf{k}}$
- e. $45 \hat{\mathbf{k}}$

22. A merry-go-round of radius $R = 2.0$ m has a moment of inertia $I = 250 \text{ kg} \cdot \text{m}^2$, and is rotating at 10 rpm. A child whose mass is 25 kg jumps onto the edge of the merry-go-round, heading directly toward the center at 6.0 m/s. The new angular speed (in rpm) of the merry-go-round is approximately


- a. 10
- b. 9.2
- c. 8.5
- d. 7.1
- e. 6.4

23. The rigid body shown is rotated about an axis perpendicular to the paper and through the point P . If $M = 0.40$ kg, $a = 30$ cm, and $b = 50$ cm, how much work is required to take the body from rest to an angular speed of 5.0 rad/s? Neglect the mass of the connecting rods and treat the masses as particles.




- a. 2.9 J
- b. 2.6 J
- c. 3.1 J
- d. 3.4 J
- e. 1.6 J

24. Two forces of magnitude 50 N, as shown in the figure below, act on a cylinder of radius 4 m and mass 6.25 kg. The cylinder, which is initially at rest, sits on a frictionless surface. After 1 second, the velocity and angular velocity of the cylinder in m/s and rad/s are respectively




- a. $v = 0; \omega = 0$.
b. $v = 0; \omega = 4$.
c. $v = 0; \omega = 8$.
d. $v = 8; \omega = 8$.
e. $v = 16; \omega = 8$.
25. A horizontal meter stick supported at the 50-cm mark has a mass of 0.50 kg hanging from it at the 20-cm mark and a 0.30 kg mass hanging from it at the 60-cm mark. Determine the position on the meter stick at which one would hang a third mass of 0.60 kg to keep the meter stick balanced.
- a. 74 cm
b. 70 cm
c. 65 cm
d. 86 cm
e. 62 cm
26. The period of a satellite circling planet Nutron is observed to be 84 s when it is in a circular orbit with a radius of 8.0×10^6 m. What is the mass of planet Nutron?
- a. 6.2×10^{28} kg
b. 5.0×10^{28} kg
c. 5.5×10^{28} kg
d. 4.3×10^{28} kg
e. 3.7×10^{28} kg



27. NOT USED

28. Three 5.0-kg masses are located at points in the xy plane in deep space. What is the magnitude of the resultant gravitational force on the mass at $x = 0, y = 0.30$ m?



- a. 2.6×10^{-8} N
- b. 2.0×10^{-8} N
- c. 2.9×10^{-8} N
- d. 2.3×10^{-8} N
- e. 2.1×10^{-8} N

1. D
2. A
3. D
4. A
5. C
6. C
7. C
8. A
9. A
10. B
11. A
12. B
13. A
14. B
15. D
16. B
17. B
18. A
19. B
20. A
21. A
22. D
23. B
24. B
25. B
26. D
27. C
28. D