


Physics at the VLHC



1. Future Colliders
2. VLHC detector issues
3. Physics Potential of the VLHC
4. Summary

Ulrich Baur

State University of New York at Buffalo

1 – Future Colliders

- e^+e^- Linear Colliders

- ☞ TESLA/NLC: $\sqrt{s} = 500 \text{ GeV} - 1.5 \text{ TeV}$

- $\mathcal{L} = \text{few} \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

- ☞ CLIC (CERN): $\sqrt{s} = 3 \text{ TeV} - 5 \text{ TeV}$

- $\mathcal{L} \approx 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$

- Muon Collider

- ☞ $\sqrt{s} = 400 \text{ GeV} - 3 \text{ TeV}$

- ☞ $\mathcal{L} = 10^{33} \text{ cm}^{-2} \text{ s}^{-1} - 5 \cdot 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

- LHC upgrade scenarios (SLHC)

studied by ATLAS (ATL-PHYS-2001-002)
and CMS:

☞ luminosity upgrade to

$$\mathcal{L} = 5 \cdot 10^{34} \text{ cm}^{-2} \text{ s}^{-1} - 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$$

☞ and/or energy upgrade: $\sqrt{s} = 28 \text{ TeV}$

requires $\sim 17 \text{ T}$ magnets (do not exist yet)

☞ **remarks**

→ for $\mathcal{L} = 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ the performance of LHC detectors is degraded, even with major upgrades (occupancy and radiation, pile-up)

→ similar problems at any hadron collider running at $\mathcal{L} \gg 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

→ in general, an increase in \sqrt{s} is easier to exploit than an increase in luminosity

- **VLHC** (Fermilab-TM-2149)

- ☞ stage 1: $\sqrt{s} = 40 \text{ TeV}$, $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

- ☞ stage 2:

\sqrt{s} (TeV)	\mathcal{L} ($\text{cm}^{-2} \text{ s}^{-1}$)
125	$5.1 \cdot 10^{34}$
150	$3.6 \cdot 10^{34}$
175	$2.7 \cdot 10^{34}$
200	$2.1 \cdot 10^{34}$

up to **50 interactions/crossing** (cf. LHC: 20)

- **remarks**

- ☞ TESLA/NLC give access to the same energy regime as the LHC. They complement the LHC

- ☞ CLIC uses a technology (two beam acceleration) very different from that used by TESLA/NLC and is a post TESLA/NLC machine

- ☞ CLIC begins to give access to energies which the LHC (without upgrade in energy) cannot reach

- remarks (cont.)

- ☞ stage 2 of the the VLHC is a post LHC and post TESLA/NLC machine

- ☞ stage 2 of the VLHC breaks completely new ground

- rest of this talk

- ☞ remarks on detector requirements

- ☞ compare LHC upgrade scenarios with stage 1 of the VLHC where appropriate

- ☞ discuss physics reach of stage 2 of the VLHC

- ☞ all estimates/extrapolations carry substantial uncertainties. More precise results should be available after Snowmass

- Result:

- regardless of what we will find at the LHC we will eventually want to have a hadron collider operating in the 100 TeV range

VLHC: UV fixed point of HEP program

2 – VLHC Detector Issues

- Physics should drive the needed detector technologies
- LHC technology should be ok for VLHC stage 1 detectors
- need serious R&D for stage 2 detectors

☞ electron detection

→ high charged track multiplicity is a potential problem

→ isolation is messy: many interactions/crossing

☞ muon detection

momentum measurement for multi-TeV μ 's is difficult and requires a very large, many Tesla magnet

☞ E_T

difficult due to many interactions/crossing

☞ jets

- need small constant term ($\sigma/E \sim 1/\sqrt{E}$)
- need to understand how many interactions/crossing influence jet energies (similar to LHC)
- need forward jet tag (up to $|\eta| = 6 - 7?$)

☞ *b*-tagging

radiation environment and track multiplicity pose problems

3 – Physics Potential of the VLHC

To illustrate the physics potential of the VLHC we consider a few more or less representative examples:

- precision SM physics and anomalous WWV ($V = \gamma, Z$) couplings
- Higgs boson physics
- supersymmetry
- strong electroweak symmetry breaking
- new gauge bosons
- compositeness (excited quarks and leptons)
- extra dimensions

Precision SM Physics

- this is not the primary reason for building the VLHC!
- well known from previous machines; many areas of the SM will have been tested at the 1-loop level
- for measurements where LHC is competitive (M_W , m_{top}), the ultimate precision is limited by systematic uncertainties. These are difficult to reduce
- special case: anomalous gauge boson couplings
 - ☞ concentrate on trilinear WWV ($V = \gamma, Z$) couplings: $\kappa_V, \lambda_V, g_1^Z$
 - ☞ non-SM contributions grow with energy ($\sim \sqrt{s}$ or $\sim s$); details depend on coupling and process considered
 - ☞ need form factor to guarantee S -matrix unitarity
 - ☞ limits depend on form factor scale Λ_{FF}
 - ☞ limits scale roughly with $(\int \mathcal{L} dt)^{1/4}$
 - increasing $\int \mathcal{L} dt$ by a factor 10, strengthens bounds by about a factor 2 – 3

- 95% CL limits:

- ☞ $\Delta\kappa_\gamma, \lambda_\gamma$ from $pp \rightarrow W(\rightarrow \ell\nu)\gamma$

- ☞ $\Delta\kappa_Z, \lambda_Z, \Delta g_1^Z$ from $pp \rightarrow WZ \rightarrow \ell_1\nu\ell_2^+\ell_2^-$

- ☞ dipole form factor (similar to proton form factor) with $\Lambda_{FF} = 10$ TeV

\sqrt{s}	14 TeV	28 TeV	40 TeV	200 TeV	CLIC (5 TeV)
$\int \mathcal{L} dt$	100 fb ⁻¹	100 fb ⁻¹	100 fb ⁻¹	200 fb ⁻¹	1 ab ⁻¹
$\Delta\kappa_\gamma$	0.034	0.027	0.023	0.013	$6 \cdot 10^{-5}$
λ_γ	0.0014	$8 \cdot 10^{-4}$	$6 \cdot 10^{-4}$	$3 \cdot 10^{-4}$	$8 \cdot 10^{-5}$
$\Delta\kappa_Z$	0.040	0.036	0.035	0.020	$7 \cdot 10^{-5}$
λ_Z	0.0028	0.0023	0.0020	0.0011	$6 \cdot 10^{-5}$
Δg_1^Z	0.0038	0.0023	0.0020	0.0011	$2 \cdot 10^{-4}$

- ☞ for larger values of Λ_{FF} , the limits improve **substantially**: $\sqrt{s} = 200$ TeV, $\Lambda_{FF} = 50$ TeV:

- $|\lambda_\gamma| < 0.0001$

- ☞ SM radiative corrections are $\mathcal{O}(\text{few} \times 10^{-4})$



- hadron and e^+e^- colliders are complementary

- ☞ hadron colliders probe high energy behaviour of helicity amplitudes


- ☞ e^+e^- colliders test angular distributions

Higgs boson physics

- the SM Higgs boson will be discovered, if it exists, at the Tevatron/LHC over the entire allowed mass range (< 1 TeV)
- measurement of SM Higgs parameters at the LHC:
 - ☞ M_H to 0.1%
 - ☞ Γ_H to $\leq 10\%$
 - ☞ $\sigma \times \text{Br}$ to 10%
 - ☞ ratios of couplings (WWH , ZZH , $\bar{t}tH$, $\bar{b}bH$) to 10 – 20%, in many cases dominated by statistics
 - ☞ weak boson fusion and forward jet tagging crucial to measure Higgs couplings
 - ☞ no information on HHH coupling (rate and/or background limited)




- precision of ratios of couplings **might** (need more studies!) improve by about a factor 2 at SLHC
- TESLA/NLC $H f \bar{f}$ couplings (**Battaglia**):



👉 similar precision for VVH ($V = W, Z$) couplings

- HHH coupling in e^+e^- collisions:



- linear colliders can measure all couplings to $\mathcal{O}(10^{-2})$
- VLHC:
 - ☞ tagging jets in weak boson fusion become more forward with increasing \sqrt{s}
 - easier to suppress bgd.
 - ☞ significantly improve LHC measurement of Higgs couplings at the VLHC (stage 1)? Measure λ_{HHH} ?
 - need detailed study

what if . . .

- no Higgs boson is found at the LHC:
 - ☞ strongly interacting Higgs sector?
 - VLHC (stage 1 may give hints already)
- Higgs boson compatible with a SM interpretation is found at the LHC, but no sparticles:
 - TESLA/NLC and/or CLIC for precision Higgs boson physics
 - VLHC for high mass sparticles search (and precision Higgs boson physics?)
- MSSM, Higgs boson(s) and some sparticles are found at the LHC:
 - CLIC and/or VLHC complete sparticle spectrum and for precision Higgs boson physics

Supersymmetry

- with 100 fb^{-1} , the LHC can find squarks (\tilde{q}) and gluinos (\tilde{g}) if their masses are $\leq 2 \text{ TeV}$
- increasing the LHC luminosity by a factor 10 extends the mass reach by about 20%.
- doubling the LHC energy to $\sqrt{s} = 28 \text{ TeV}$ provides access to \tilde{q} and \tilde{g} with masses up to 3–4 TeV
→ at stage 1 of the VLHC one can detect squarks and gluinos with masses up to 4 – 5.5 TeV
- LHC: other sparticles are mainly detected from \tilde{q} and \tilde{g} cascade decays
☞ for many mSUGRA models, the LHC will miss most of the sleptons, charginos and neutralinos, and the heavy Higgs bosons
- one can construct inverted hierarchy models (IHM) where none of the sparticles can be discovered (5σ) at the LHC (Baer et al.)

- stage 2 of the VLHC might be able to probe the dynamics of SUSY breaking
 - ☞ any SUSY theory must contain a mechanism for breaking SUSY
 - ☞ and a method (messengers) for communicating SUSY breaking to the sparticles
 - ☞ two scales:
 - SUSY breaking vev F
 - messenger scale M
 - ☞ sparticle mass:

$$\tilde{m} \sim \eta \frac{F}{M}$$

η : dimensionless suppression factor from coupling constants

- ☞ GMSB: M is replaced by vector-like messenger field; $\eta \sim \alpha/4\pi$
 - for $\sqrt{F} \sim M$, both messenger fields and SUSY breaking scale could be as low as **10 – 100 TeV**
 - **could be accessible at stage 2 of the VLHC**

☞ M can be measured from sparticle spectroscopy

→ expected precision at the LHC: $\sim 30\%$

☞ F from NLSP lifetime and mass

- SUSY mass scales:

☞ electroweak scale protected if superpartners coupling most strongly to Higgs boson have masses < 1 TeV

☞ \tilde{t}, \tilde{b}_L , weak gauginos, higgsinos have $m < 1$ TeV

☞ other squarks/sleptons contribute to weak scale at two loop


→ $m < 20$ TeV

what if . . .

- the LHC finds \tilde{q} and \tilde{g} and maybe a few other sparticles
 - VLHC and CLIC have a good chance to fill in the gaps of the sparticle spectrum
- the LHC finds \tilde{t} and \tilde{g} but misses the first two generation squarks
 - VLHC (maybe stage 1, but certainly stage 2) should find the missing squarks (no quantitative estimates so far)
- the LHC discovers SUSY and finds it is low energy GMSB
 - stage 2 of the VLHC can probe messenger sector

strong electroweak symmetry breaking

- if no Higgs boson exists, one expects that longitudinal W 's and Z 's interact strongly for $\sqrt{\hat{s}} \geq 1$ TeV
- vector boson scattering, eg:




- forward jet tagging and central jet veto are powerful tools to reduce background
- **example:**
 - ☞ non-resonant scattering
 - ☞ most difficult case
 - ☞ best channel: $W^\pm W^\pm \rightarrow \ell_1^\pm \nu \ell_2^\pm \nu$
 - ☞ compare **1 TeV SM Higgs boson** with **K -matrix unitarization model** (Bagger et al.)


☞ K -matrix unitarization: replace partial wave amplitudes a_l^I by

$$t_l^I = \frac{a_l^I}{1 - ia_l^I}$$

- significance versus \sqrt{s} :



- signal and background have same shape
 - large statistics needed for a convincing signal
- (ATLAS)



- ☞ hatched: WW and WZ background
- ☞ solid: K -matrix unitarization
- ☞ dashed: 1 TeV SM Higgs boson


- LHC, at $\mathcal{L} = 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$:
 - ☞ degradation of forward jet tag and central jet veto due to pile-up
 - ☞ large ($\approx 50\%$) probability for fake jet tags even at momenta of a few hundred GeV
 - luminosities $\mathcal{L} > 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ do not help much

what if . . .

- LHC does not find a Higgs boson but observes hints for strong electroweak symmetry breaking
 - stage 1 of the VLHC should find convincing signal
 - fully explore strong dynamics at stage 2 of the VLHC

Extra gauge bosons

- additional gauge bosons, W' and Z' , appear in many GUT models (E_6, \dots)
- the reach depends on the W' , Z' couplings to quarks and charged leptons
- concentrate on $Z' \rightarrow \mu^+ \mu^-$ with SM couplings here (classic benchmark)



👉 similar reach for W' 's

- can measure:
 - ☞ Z' mass at (energy or luminosity upgraded) LHC to $< 1\%$
 - ☞ Z' width to a few percent
 - ☞ couplings using other channels: $Z' \rightarrow jj$, $Z' \rightarrow W^+W^-$
- CLIC: from **indirect measurements**: sensitivity up to $M_{Z'} = 30$ TeV
- **direct** search at CLIC: only for $M_{Z'} < \sqrt{s}$
 - ☞ can measure Z' mass to $< 10^{-4}$
 - ☞ Z' width and peak cross section to better than 1%

Compositeness

- if quarks and/or leptons are composite with a scale Λ (scale of interactions which binds constituents):
 - ☞ for $\sqrt{\hat{s}} \ll \Lambda$: contact interactions
 - ☞ for $\sqrt{\hat{s}} \geq \Lambda$: production of excited quarks (q^*) and leptons (ℓ^*)
- **contact interactions**: example: 2-jet events
 - ☞ expect excess of high E_T centrally produced jets
 - ☞ maximum scale probed for 300 fb^{-1} :

$\sqrt{\hat{s}}$ (TeV)	Λ (TeV)
14	40
28	60
40	~ 75
100	~ 115
200	~ 130

- excited quarks:

- ☞ produced via qg fusion in s -channel: $qg \rightarrow q^*$

- ☞ decays: $q^* \rightarrow qg, q\gamma, qW, qZ$

- ☞ effective Lagrangian for $q^*q\gamma$ coupling is of magnetic moment type


$$\mathcal{L} \sim \frac{f_s g}{\Lambda} q^* \sigma_{\mu\nu} F^{\mu\nu} q$$

- mass reach for $q^* \rightarrow jj, f_s = 1, M_{q^*} = \Lambda$:

- ☞ LHC, 100 fb^{-1} (1000 fb^{-1}): **7 TeV** (**8 TeV**)

- ☞ $\sqrt{s} = 28 \text{ TeV}$, 100 fb^{-1} (1000 fb^{-1}): **10 TeV** (**11 TeV**)

👉 VLHC: 5σ reach for $f_s = 1$, $M_{q^*} = \Lambda$:




👉 for $f_s = 0.1$ the reach is about a factor 2 smaller

what if ...


- the LHC finds evidence for contact interactions?
 - $\Lambda < 60$ TeV
 - find excited quarks and/or leptons at the VLHC (stage 2)

Extra dimensions

- Fields propagating in more than 4 dimensions lead to Kaluza-Klein (KK) excitations, modifications to cross section, or E_T signatures
- example: $jet+graviton$ production in ADD model; graviton manifests as E_T
 - ☞ cross section depends on M_D , the scale of gravity and δ , the number of extra dimensions ($\delta = 1$ ruled out by celestial mechanics)
 - ☞ main background: $Z(\rightarrow \bar{\nu}\nu) + jets$ (Hinchliffe)




- M_D reach of the (upgraded) LHC and VLHC:



- warped extra dimensions (RS models):
 - ☞ SM gauge and fermion fields live on the TeV-brane
 - ☞ or they may propagate in the bulk
- SM fields constrained to the TeV brane:
 - ☞ colliders are KK resonance factories
 - ☞ production of graviton KK excitations ($G^{(n)}$):
 $\bar{q}q, gg \rightarrow G^{(n)} \rightarrow \ell^+ \ell^-$

- example: LHC (Davoudiasl et al.)




👉 red: $m_{G^{(1)}} = 1 \text{ TeV}$, $k/\bar{M}_{Pl} = 0.1$

👉 green: $m_{G^{(1)}} = 1.5 \text{ TeV}$, $k/\bar{M}_{Pl} = 0.2$

➔ k : AdS_5 curvature scale

- the LHC with 100 fb^{-1} can determine the spin-2 nature of a KK graviton for $m_{G^{(1)}} \leq 4.2 \text{ TeV}$
 - ➔ no VLHC studies yet (after Snowmass?)
 - 👉 CLIC: sensitive to $G^{(1)}$ up to kinematic limit

- if no evidence for new particles at LHC or TESLA/NLC:
 - ☞ search for indirect effects of KK excitations through contact like interactions
 - ☞ example (SM fields propagating in the bulk) 95% CL, Drell-Yan production (Davoudiasl et al.):



- ☞ m_1^{\min} : mass of lightest KK excitation
- ☞ ν : bulk mass parameter; controls how far off the TeV-brane ($\nu \rightarrow \infty$) the wave function is located
- ☞ black: Tevatron, Run I, red: Tevatron, 2 fb^{-1} , blue: Tevatron, 30 fb^{-1} , green: LHC, 10 fb^{-1} , cyan: LHC, 100 fb^{-1}
- ➔ no VLHC studies yet (after Snowmass?)

what if . . .

- the LHC finds evidence for extra-dimensions?
 - the VLHC (stage 2) will directly probe M_D
 - VLHC could find totally unexpected physics

VLHC Pocket Guide

channel	LHC	LHC	28 TeV	40 TeV	200 TeV
particle	100 fb^{-1}	1 ab^{-1}	100 fb^{-1}	100 fb^{-1}	100 fb^{-1}
\tilde{q}, \tilde{g}	2	2.5	4	5.5	> 10
$W' Z'$	4.5	5.4	7	8.5	33
q^*	7	8	10	13	50
$\Lambda \text{ comp.}$	33	50	60	75	130
$M_D (\delta = 2)$	9	12	15	20	75

- large uncertainties
- not exhaustive
- all masses in TeV

4 – Summary

- Upgrading the LHC luminosity by a factor 10 increases the reach by 20%
- Doubling the LHC energy to $\sqrt{s} = 28 \text{ TeV}$ increases the reach by up to a factor 2
- stage 1 of the VLHC only insignificantly increases the reach of a 28 TeV LHC
 - makes only sense if LHC is not significantly upgraded in energy
- At some point we will, inevitably, want to go to the 100 TeV region
- the VLHC is the only machine which can directly discover new physics in the multi 10 TeV region
- most of the what if . . . scenarios discussed suggest that we need the VLHC
 - we don't need to wait for LHC results to decide
- need a coordinated and coherent international plan for the VLHC which is part of a comprehensive and global HEP program