

#0) (One more from Ch.6)

If the generalized driving forces  $Q_i$  are not sinusoidal, show that the forced vibrations of the normal coordinates in the absence of damping are given by

$$\zeta_i = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \frac{G_i(\omega)}{\omega_i^2 - \omega^2} e^{-i\omega t} d\omega,$$

where  $G_i(\omega)$  is the Fourier transform of  $Q_i$  defined by

$$Q_i(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} G_i(\omega) e^{-i\omega t} d\omega.$$

If the dissipation function is simultaneously diagonalized along with  $T$  and  $V$ , show that the forced vibrations are given by

$$\zeta_i = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \frac{G_i(\omega)(\omega_i^2 - \omega^2 + i\omega\mathcal{F}_i)}{(\omega_i^2 - \omega^2)^2 + \omega^2\mathcal{F}_i^2} e^{-i\omega t} dt,$$

Physics 6321:  
Homework #9:  
Prof. Olness  
Spring 2024

---

**PROBLEM #1:** Consider the metric:

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and vector:

$$v = \begin{pmatrix} x \\ y \end{pmatrix}$$

- a) Compute the length-squared of the vector  $v$ .
- b) Write down a rotation matrix  $R$ .
- c) Show the length of  $v$  is invariant under a rotation.
- d) Using Mathematica, make a ContourPlot of the length of the vector in the  $\{x,y\}$  space. Comment about the result.

**PROBLEM #2:** Consider the metric:

$$g_{\mu\nu} = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

and vector:

$$v = \begin{pmatrix} x \\ y \end{pmatrix}$$

- a) Compute the length-squared of the vector  $v$ .
- b) Using Mathematica, make a ContourPlot of the length of the vector in the  $\{x,y\}$  space. Comment about the result.
- c) Can you think of a physical example where this is the appropriate metric to use to measure distance?? (There are more than one answer, but the more natural the better.)

**PROBLEM #3:** Consider the metric:

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

and vector:

$$v = \begin{pmatrix} t \\ x \end{pmatrix}$$

- a) Compute the length-squared of the vector  $v$ .
- b1) Write down a boost matrix  $B$  in terms of  $\beta$  and  $\gamma$ .
- c1) Show the length of  $v$  is invariant under a boost.
- b1) Write down a boost matrix  $B$  in terms of  $\cosh$  and  $\sinh$ .
- c1) Show the length of  $v$  is invariant under a boost in terms of  $\cosh$  and  $\sinh$ .
- d) Using Mathematica, make a ContourPlot of the length of the vector in the  $\{t,x\}$  space. Comment about the result.

**PROBLEM #4:** For the general mass case of  $p_1 + p_2 \rightarrow p_{12}$  compute the components of all 4-vectors in terms of invariants of the problem:  $\{m_1^2, m_2^2, s\}$ . Assume the 3-momentum lies along the  $z$ -axis. Hint: The  $z$ -component of the momentum should be proportional to:  $\Delta(s, m_1^2, m_2^2)$ , where  $\Delta(a, b, c) = \sqrt{a^2 + b^2 + c^2 - 2(ab + bc + ca)}$ .

**PROBLEM #5:** For the zero mass case of  $p_1 + p_2 \rightarrow p_3 + p_4$  compute the components of all 4-vectors in terms of invariants of the problem:  $\{s, t, u\}$ . Assume the 3-momentum of  $\{p_1, p_2\}$  lies along the  $z$ -axis.

**PROBLEM #6:** Consider the reaction:  $p\bar{p} \rightarrow 3X$  where  $X$  has a rest mass of  $140\text{GeV}$ . Compute the threshold beam energy for

- a) colliding beams, and
- b) for a fixed target experiment.

**PROBLEM #7:** Consider the rapidity  $y$  and the pseudo-rapidity  $\eta$ :

$$y = \frac{1}{2} \ln \left( \frac{E + P_L}{E - P_L} \right)$$

$$\eta = -\ln \left[ \tan \left( \frac{\theta}{2} \right) \right]$$

- Make a parametric plot of  $\{y, \eta\}$  as a function of  $m/E$  where  $m$  is the mass of the particle.
- Show that in the limit  $m \rightarrow 0$  that  $y \rightarrow \eta$ .
- Make a table of  $\eta$  for  $\theta = [0^\circ, 180^\circ]$  in steps of 5 degrees.
- Make a table of  $\theta$  for  $\eta = [0, 10]$  in steps of 1.

**PROBLEM #8:** For one-particle phase space, show the following equality (*with all the steps*):

$$\frac{d^3\vec{P}}{(2\pi)^3 2E} = (2\pi)\delta(P^2 - m^2) \frac{d^4P}{(2\pi)^4}$$

**PROBLEM #9:** In a Tevatron detector, consider two particles traveling in the transverse direction:

$$p_1^\mu = \{E, 100, 0, 1\}$$

$$p_2^\mu = \{E, 100, 1, 0\}$$

where the components are expressed in GeV units.  $E$  is defined such that the particles are massless.

- Compute  $E$ .
- For each particle, compute the pseudorapidity  $\eta$  and azimuthal angle  $\phi$ .
- Explain how the above exercise justifies the correct jet radius definition to be:

$$R = \sqrt{\eta^2 + \phi^2}$$

In particular, why is the above correct and  $R = \sqrt{\eta^2 + 2\phi^2}$ , for example, incorrect.

**PROBLEM #10:** Consider the reaction:  $pp \rightarrow pp$  ( $12 \rightarrow 34$ ) with **CMS** scattering angle  $\theta$ . The CMS energy is  $\sqrt{s} = 2 TeV$ .

- Compute the boost from the CMS frame to the rest frame of #2 (lab frame)
- Compute the energy of #1 in the lab frame.
- Compute the scattering angle  $\theta_{lab}$  as a function of the CMS  $\theta$  and invariants.

**PROBLEM #11:** Write the  $4 \times 4$  EM Field Strength tensor  $F^{\mu\nu}$  in terms of  $\{E, B\}$  fields.

Now apply a boost (Lorentz transformation) along the  $z$  axis and compute the transformed  $F^{\mu\nu}$ .

Examine how the  $\{E, B\}$  fields transform and check against your EM textbook results.