

-
-
8. Derive the electric field for a point charge at the origin $\left(\vec{E} = k\frac{q\vec{r}}{r^3}\right)$ from Gauss' law.
9. Solve Jackson Chapter 1 - Problem 1. Use Gauss' theorem (and $\oint \vec{E}\cdot d\vec{l} = 0$ if necessary) to prove the following:
- (a) Any excess charge placed on a conductor must lie entirely on its surface.
 - (b) A closed, hollow conductor shields its interior from fields due to charges outside, but does not shield its exterior from fields due to charges placed inside it.
 - (c) The electric field at the surface of a conductor is normal to the surface and has a magnitude σ/ϵ_0 , where σ is the charge density per unit area on the surface.
10. Consider a spherically symmetric charge distribution, i.e. $\rho(\vec{r}) = f(r)$.
- (a) Use Gauss' law to obtain an expression for the electric field as an integral of $f(r)$.
 - (b) Find the potential as a **double** radial integral with the requirement that $\Phi(r) \rightarrow 0$ as $r \rightarrow \infty$.
 - (c) Find the electric field and potential for a charge density

$$\rho(\vec{r}) = Q\delta^3(\vec{r}) - \frac{Q}{8\pi r_0^3}e^{-r/r_0} .$$

- (d) What is the total charge of the system?
- (e) Is your answer for $\Phi(r)$ what you would expect from a multipole expansion?
Comment.